



THE THREE SHIRES MEDICAL PRACTICE

Colerne, Marshfield, Pucklechurch and Wick Surgeries

Dr Richard Greenway
Dr Richard Prince
Dr Pedro Pinto
Dr Keira Prince
Dr Dana Parr

Colerne Surgery Tel: (01225) 742028
Marshfield Surgery Tel: (01225) 891265
Pucklechurch Surgery Tel: (0117) 937 2350
Wick Surgery Tel: (0117) 937 2214

HYPERTENSION (High Blood Pressure)

This leaflet explains what impacts having high blood pressure may have upon your health, and what you can do to address it.

What is blood pressure?

Blood pressure is a measure of the pressure your heart creates. As it pumps, it contracts (systolic) and relaxes (diastolic). In adults, systolic pressure is about 120 and diastolic pressure 70, written as 120/70. However, just as there is no 'normal' height, there is no 'normal' blood pressure.



So, what is hypertension?

Hypertension means high blood pressure, i.e. a reading that is higher than considered 'normal' for you. A reading of about 140/90 or over is usually considered to be high.

Who is at risk of having hypertension?

10-20% of UK adults have hypertension. Men are more at risk, and it is more common in older adults.

What causes hypertension?

Blood pressure rises when there is increased resistance to the heart pumping. Usually, there is no obvious cause; this is called 'essential hypertension' and seems to run in some families. In less than 1 in 20, there is an underlying cause, such as a kidney disorder. Temporary hypertension can occur in pregnancy.

Is hypertension dangerous?

Hypertension itself is not dangerous but can increase the risk of having a heart attack or stroke in the future. In a middle-aged man, for example, each rise of 10mm systolic pressure increases his risk of a heart attack or stroke by 20%.

What are the symptoms?

People with hypertension usually feel perfectly well. Severe hypertension may cause giddiness, breathlessness, visual problems, or headaches.



What can help?


Lifestyle change is important (see below) and may be enough to keep your blood pressure at a normal level. Your doctor may also consider drug treatment. You may need to try different sorts and doses before you get the right combination for you. They can include:

- **Water tablets (diuretics) e.g. Indapamide** – increase the amount of salt that your kidneys remove.
- **Beta-blockers e.g. Bisoprolol** – make the heart beat less strongly. Can cause tiredness. Often not suitable for asthmatics.
- **ACE inhibitors e.g. Ramipril** – relax the blood vessels. Side effects are rare with the occasional dry cough. A kidney blood test is important.
- **Calcium antagonists e.g. Amlodipine** – relax the blood vessels. Side effects can include swollen ankles.
- **Alpha-blockers e.g. Doxazosin** – relax blood vessels. They may cause a stuffy nose.

What can you do?

There are several areas in which you can make changes, in order to try and lower your blood pressure. These are known as 'lifestyle changes', and should include:

- Manage your weight.
- Eat less salt – salt influences your blood pressure.
- Eat more potassium – found in fresh fruit and vegetables.
- Take more exercise – try to have a brisk 20-minute walk each day.
- Reduce the stress in your life. Stress does not cause high blood pressure but can aggravate it.
- Stop smoking. (Advice and support available from your GP surgery)
- Reduce your alcohol intake (Less than 14 units per week, with a few alcohol-free days.)



If you need more support around your high blood pressure or have any questions, please book an appointment with a clinician at your surgery